

[GNFAC Avalanche Forecast for Thu Feb 27, 2020](#)

Good Morning. This is Doug Chabot with the Gallatin National Forest Avalanche Forecast on Thursday, February 27th at 7:15 a.m. Today's forecast is sponsored by [Spark R&D](#) and [Montana Alpine Guides](#). This forecast does not apply to operating ski areas.

Mountain Weather

In the last 24 hours, Cooke City picked up 3" of new snow with a trace to 1" falling everywhere else. Under clearing skies mountain temperatures are in the teens and wind is blowing W-NW at 10-20 mph and gusting to 25-40 mph. Today will be sunny and temperatures will climb into the 30sF. Wind will continue to be westerly and gusty. Expect another sunny day tomorrow and snow this weekend.

Snowpack and Avalanche Discussion



Bridger Range Northern Gallatin Northern Madison

Yesterday, wind drifted snow at the ridgelines. Overall these drifts were not very sensitive. However, a cornice broke midday atop the east-facing Hourglass Chute north of Bridger Bowl and triggered a 3-4 foot deep x 100-foot wide slide ([photos and details](#)). The avalanche broke on sugary facets at the ground and serves as a reminder that this layer is still a concern throughout the northern mountains. Dave rode into Wheeler Mountain in the northern Gallatin Range yesterday and looked at old crowns from a few weeks ago where he found the same layer breaking clean in his snowpit, a sign of lingering instability ([video](#)). This also mirrored his finding from Beehive Basin on Monday ([video](#)). Yesterday, skiers in Hyalite found 2' thick wind drifts on Maid of the Mist which were glued down, a positive sign, but not an ironclad one. They did not get any shooting cracks, collapses or other red flags. These would be signs to avoid avalanche terrain. Winds continued last night and triggering fresh wind-drifts remains possible, which, if you were unlucky, could break deep into old snow. For today, the avalanche danger is rated **MODERATE** on all slopes.



Southern
Madison

Southern
Gallatin

Lionhead
Range

Cooke
City

Island
Park

The mountains around Cooke City received 3" of snow without much wind. Like the southern Gallatin Range, southern Madison Range and Lionhead area, the snowpack is similarly stable. A few rogue wind slabs might be triggered, but these would be small. Ian investigated the snowpack both south and north of Cooke City and reported good stability ([video](#)). Sugary facets at the ground remain an afterthought since it would take a large load or getting extremely unlucky to trigger a slide. Around Lionhead and the other southern ranges there has been no recent avalanches or red flags ([video](#)). This zone has not gotten much snow lately and its good stability reflects that lack of loading. In general, triggering avalanches is unlikely and the danger is rated **LOW**.

If you get out, please send us your observations no matter how brief. You can fill out an [observation form](#), email us (mtavalanche@gmail.com), leave a VM at 406-587-6984, or Instagram ([#gnfacobs](#)).

Upcoming Avalanche Education and Events

Our education calendar is full of awareness lectures and field courses. Check it out and plan to attend one or two: [Events and Education Calendar](#).

COOKE CITY

Every Friday and Saturday, **Snowpack Update and Rescue Training**. Friday, 6:30-7:30 p.m. at the Soda Butte Lodge. Saturday anytime between 10-2 @ Round Lake.

BOZEMAN

February 27 - March 1, **Bozeman Splitfest**, More [info and schedule here](#).

March 4, **1-hr Avalanche Awareness**. 6-7 p.m. at REI.

BILLINGS

March 12, **1-hr Avalanche Awareness**, 6-7 p.m. at The Basecamp.